

Protocols

Clinical Opiate Withdrawal Scale (COWS)

Flowsheet for measuring symptoms over a period of time during buprenorphine induction.

For each item, write in the number that best describes the patient's signs or symptom. Rate on just the apparent relationship to opiate withdrawal. For example, if heart rate is increased because the patient was jogging just prior to assessment, the increase pulse rate would not add to the score.

Patient's Name: _____ Date: _____				
Buprenorphine induction:				
Enter scores at time zero, 1 hour after first dose, at each new dose, 1 hour after every dose				
Times: _____				
Resting Pulse Rate: (record beats per minute) <i>Measured after patient is sitting or lying for one minute</i> 0 pulse rate 80 or below 1 pulse rate 81-100 2 pulse rate 101-120 4 pulse rate greater than 120				
Sweating: <i>over past ½ hour not accounted for by room temperature or patient activity</i> 0 no report of chills or flushing 1 subjective report of chills or flushing 2 flushed or observable moistness on face 3 beads of sweat on brow or face 4 sweat streaming off face				
Restlessness <i>Observation during assessment</i> 0 able to sit still 1 reports difficulty sitting still, but is able to do so 3 frequent shifting or extraneous movements of legs/arms 5 unable to sit still for more than a few seconds				
Pupil size 0 pupils pinned or normal size for room light 1 pupils possibly larger than normal for room light 2 pupils moderately dilated 5 pupils so dilated that only the rim of the iris is visible				
Bone or Joint aches <i>If patient was having pain previously, only the additional component attributed to opiates withdrawal is scored</i> 0 not present 1 mild diffuse discomfort 2 patient reports severe diffuse aching of joints/muscles 4 patient is rubbing joints or muscles and is unable to sit still because of discomfort				

PATIENT LABEL



XOPWDSCALE

Summa Health System

90281125 (11/19)

Clinical Opiate Withdrawal Scale (COWS), continued

Runny nose or tearing <i>Not accounted for by cold symptoms or allergies</i> 0 not present 1 nasal stuffiness or unusually moist eyes 2 nose running or tearing 4 nose constantly running or tears streaming down cheeks				
GI Upset: <i>over last 1/2 hour</i> 0 no GI symptoms 1 stomach cramps 2 nausea or loose stool 3 vomiting or diarrhea 4 multiple episodes of diarrhea or vomiting				
Tremor <i>observation of outstretched hands</i> 0 no tremor 1 tremor can be felt, but not observed 2 slight tremor observable 4 gross tremor or muscle twitching				
Yawning <i>observation during assessment</i> 0 no yawning 1 yawning once or twice during assessment 2 yawning three or more times during assessment 4 yawning several times/minute				
Anxiety or irritability 0 none 1 patient reports increasing irritability or anxiousness 2 patient obviously irritable or anxious 4 patient so irritable or anxious that participation in the assessment is difficult				
Gooseflesh skin 0 skin is smooth 3 piloerection of skin can be felt or hairs standing up on arms 5 prominent piloerection				
Total scores with observer's initials				

Score:

5-12 = mild;

13-24 = moderate;

25-36 = moderately severe;

more than 36 = severe withdrawal

Notify Obstetric resident for scores > 24